

A decade of canopy-tree seedling survival and growth in two Bornean rain forests: persistence and recovery from suppression

LISA J. DELISSIO^{*1}, RICHARD B. PRIMACK[†], PAMELA HALL[‡] and H. S. LEE[§]

^{*}*Salem State College, Department of Biology, 352 Lafayette Street, Salem, MA 01970, USA*

[†]*Boston University, Department of Biology, Boston, MA 02215, USA*

[‡]*Aarhus University Herbarium, Building 137, Universitetsparken, DK-8000 Aarhus C, Denmark*

[§]*Forest Department, 93660 Kuching, Malaysia*

(Accepted 3rd September 2001)

ABSTRACT. The population dynamics of 8500 shade-tolerant tree seedlings of 13 tree species were followed for 10 years at rain forests in Lambir Hills National Park and Bako National Park, Malaysian Borneo. Since these dipterocarp forests have lower rates of canopy gap formation than do rain forests elsewhere, tree seedling biology was predicted to differ. Approximately 50% of seedlings present in 1986 were still alive in 1996. Seven out of thirteen had seedling populations composed predominantly of individuals that were at least 10 years old. These seedlings can undergo alternating periods of relatively rapid and slow growth. Many seedlings that grew rapidly in the first census interval survived through a period of suppression in the second census interval, and conversely many seedlings with a history of suppression had exceptionally high growth in the final census interval. Seedlings of South-East Asian forest tree species are long-lived and appear to be adapted to long periods of suppression in the understorey.

KEY WORDS: age structure, dipterocarps, growth, Lambir Hills National Park, mortality, population dynamics, resilience, Sarawak, shade-tolerant tree seedlings, tropical rain forest

INTRODUCTION

In the species-rich, aseasonal, lowland forests of Borneo, shade-tolerant trees of the Dipterocarpaceae, the dominant tree family, as well as canopy trees of several other families, display gregarious flowering events at intervals of approximately 2–11 y (Appanah 1985, Ashton 1988, Curran *et al.* 1999, Wich & van Schaik 2000, Wood 1956). Mast-fruiting, the supra-annual production of

¹ Corresponding author. Email: lisa.delissio@salemstate.edu

fruits, follows. This masting yields large numbers of seeds, up to 115 000 per individual dipterocarp (Ashton 1988). The seeds that survive germinate within weeks resulting in dense seedling populations with clumped distributions (Ashton 1969, Curran & Webb 2000, Fox 1967, Itoh 1995). Seedling mortality rates tend to be high initially, decreasing over 3–4 y (Barnard 1956, Curran & Webb 2000, Delissio 2000, Itoh *et al.* 1995, Wyatt-Smith 1958).

Most shade-tolerant tree seedlings require light from canopy gaps for survival and growth into canopy-sized trees (Platt & Strong 1989, Swaine & Whitmore 1988). For example, at La Selva tropical rain forest in Costa Rica, approximately 75% of canopy tree species must experience canopy gaps in order to reach maturity (Hartshorn 1978). Periodic large-scale disturbances such as hurricanes and cyclones, earthquakes, fires and volcanism are important determinants of forest dynamics and structure in the rain forests of the Caribbean region, Panama, Hawaii, New Guinea, Australia, the Solomon Islands, and many parts of South-East Asia, including northern and eastern Borneo (Adams 1992, Arriaga 2000, Beaman *et al.* 1985, Brokaw & Walker 1991, Clark 1990, Crow 1980, Garwood *et al.* 1979, Putz & Milton 1982, Vitousek *et al.* 1993, Walker 1991, Whitmore 1989, 1998). These disturbances result in snapped trunks and uprooted trees which create gaps in the forest canopy that allow light to penetrate into the understorey.

In contrast to other South-East Asian or neotropical forests, the primary rain forests of Sarawak in north-west Borneo have a lower incidence of large-scale natural disturbances. While fires and dry periods do occur (Baillie 1976, Brünig 1969), extended droughts are rare (Nakagawa *et al.* 2000, Walsh 1996). Trees in these forests are more likely to die standing and fall apart in place (Ashton & Hall 1992, Gale 2000, Hall 1991), creating smaller, shorter-lived gaps (Whitmore 1978). Moreover, in at least one of two forests considered in this investigation (Lambir Hills National Park), gaps form at a relatively slow rate, 0.1% of the total area y^{-1} (Ohkubo *et al.* submitted), compared with rates of 0.7–2.9% y^{-1} in neotropical forests and in the forests of French Guiana (Clark 1990, van der Meer 1995). The most important form of disturbance in Lambir is likely to be the rare catastrophic landslides that occur on steep slopes as a result of heavy rains (Ohkubo *et al.* 1995).

The relative rarity of treefall events in the forests of north-west Borneo, and the subsequent rarity of canopy gaps, may have important implications for the evolution of life history characteristics in shade-tolerant tree species, and for the maintenance of high species diversity in these forests. Moreover, the lowland dipterocarp forests of South-East Asia are of interest to evolutionary ecologists because of their unusually high numbers of coexisting, closely related tree species (Ashton 1969, Whitmore 1984, 1995) which also form the basis of the South-East Asian timber industry. Therefore, the information from long-term census work has important ecological and economic implications.

METHODS

Study sites

The research sites for our study are located within two large forest reserves in the Malaysian state of Sarawak: Lambir Hills National Park and Bako National Park. In 1965, Ashton (Ashton & Hall 1992), established these sites for an on-going long-term study of mature trees. Each site has four permanent 0.6-ha plots numbered 1–4 at Lambir and 2–5 at Bako. The plots at Lambir are situated on a mosaic of sandy humult ultisols to udult clay ultisols (Ashton & Hall 1992, Palmiotto 1998). The plots at Bako are on sandy humult ultisols (Ashton & Hall 1992, Hall 1991). The research plots at Lambir and Bako span altitudinal ranges of 360–450 m asl, and 30–75 m asl respectively (Hall 1991).

The research plots were established in mature forest in 1965. Data collected from these plots over the 1965–1985 period were used to analyse forest structure and community composition (Ashton & Hall 1992, Hall 1991, Primack & Hall 1992). Since 1965, they have accumulated gaps (Ashton & Hall 1992). The plots at Bako appear to be recovering from a large-scale disturbance, perhaps a drought (Hall 1991, Primack & Hall 1992), and Plots 3 and 4 appear to be recovering from more recent smaller-scale disturbances (L. Delissio, *pers. obs.*). Bako is a relatively low-stature forest with a high density of smaller trees, and the number of smaller trees is declining as the canopy increases in biomass. The forest at Lambir is of higher stature, and is relatively stable in stand structure compared with Bako. No large-scale disturbances occurred in the research plots during this study.

The Lambir plots are now embedded in a 52-ha long-term research plot, established for the investigation of tree demographics and reproductive ecology (Davies *et al.* 1998, Itoh *et al.* 1997, Lee *et al.* 1995, Yamakura *et al.* 1995). For trees ≥ 10 cm in diameter at breast height (dbh), this forest is among the most species-rich forests in the world (Condit 1995).

Study species

The 13 study species (seven at Bako and six at Lambir) selected in 1986, are shade-tolerant climax species as defined by Swaine & Whitmore (1988). Four dipterocarp species were included because of their importance to the ecology of these forests. The following criteria were applied to select species: (1) trees of the species were common within the site; (2) seedlings of the species could be identified in the field; and (3) the species was of ecological or silvicultural importance (Hall & Primack 1987).

The focal species at Bako included *Artocarpus odoratissimus* Blanco, *Cotylelobium melanoxyton* (Hook. f.) Pierre, *Dryobalanops beccarii* Dyer, *Gonocaryum minus* Sleum., *Pimelodendron griffithianum* (Muell. Arg.) Benth. and *Swintonia schwenkii* (Teijsm. & Binn.) Teijsm. & Binn. (Table 1). The focal species at Lambir

included *Allantospermum borneensis* Form., *Dipterocarpus globosus* Vesq., *Elateriospermum tapos* Bl., *Mangifera parvifolia* Boerl. & Koord., *Mangifera pentandra* Hook., *Shorea beccariana* Burck, and *Whiteodendron moultonianum* (W.W. Sm.) Steen. (Table 1). Fieldwork showed that some *S. beccariana* juveniles are morphologically indistinguishable from those of *Shorea amplexicaulis* Ashton (see Ashton 1969). The two species are combined in this analysis and referred to as *Shorea beccariana*. In previous reports, *M. parvifolia* was referred to by its synonym, *Mangifera havilandii*.

Census

In six 100-m × 60-m plots (three at Lambir and three at Bako), four 2-m × 100-m seedling transects spaced 10 m apart were established (Hall & Primack 1987). Two plots that were irregular in shape (Plot 4 which runs along a ridge at Lambir and Plot 2 at Bako) had a different arrangement of 2-m-wide transects, but the same total transect area as the 100-m × 60-m plots. Each transect was divided into 10-m sections marked by permanent pegs. Both sites had transects that were offset to avoid trails. The total census area at Lambir was 3200 m². At Bako, some transects were shortened to avoid trails for a total census area of 3000 m². All eight plots were sampled at each of the four censuses (1986, 1988, 1990 and 1996).

Seedlings were defined as individuals with a main stem < 1 cm dbh and ≥ 20 cm in height. All seedlings of the study species within the transects were censused. Each seedling was mapped and labelled with a permanent nursery tag. Seedlings within the sampling area that had recently attained 20 cm were added at each census. Positive growth was recorded, as was lost height due to dieback and breakage. *Dryobalanops beccarii* recruits only were subsampled within three of the plots at Bako in 1996 due to their large numbers. The total area sampled for *D. beccarii* recruits in 1996 was 1970 m².

Analysis

Growth rates were calculated as the difference in height divided by the difference in the decimal dates of the censuses. Median and maximum annualized growth rates (cm y⁻¹) were calculated for seedlings present in 1986 that survived to 1996 and that had a net positive change of height over that interval. The proportion of seedlings surviving was calculated as the difference between the number of seedlings present in 1986 and the number of this set that had died as of 1996 divided by the number of seedlings present in 1986. Seedlings that had advanced to the sapling size class (> 1 cm dbh) were counted as surviving.

RESULTS

At the 1986 census 4622 seedlings were tagged. Of this initial set, 2373 were alive and seedling-sized at the 1996 census. For all species from both sites

Table 1. Study species, sample sizes and population densities of seedlings of each species present at the 1986 census, and sample sizes and population density of recruits of each species at the census in which they were initially tagged (1988, 1990 or 1996). Most species did not occur in all plots.

Species/Family		N	Population density (seedlings per m ²) by plot				
			2	3	4	5	
Lambir							
<i>Allantospermum borneensis</i> Ixonanthaceae	Recruits:	1986	205	—	0.070	0.158	0.029
		1988	18	—	0.010	0.008	0.006
		1990	3	—	0.000	0.001	0.003
		1996	301	—	0.150	0.093	0.033
<i>Dipterocarpus globosus</i> Dipterocarpaceae	Recruits:	1986	1881	—	0.334	1.095	0.923
		1988	105	—	0.026	0.04	0.056
		1990	22	—	0.001	0.018	0.009
		1996	51	—	0.014	0.026	0.009
<i>Elatiospermum tapos</i> Euphorbiaceae	Recruits:	1986	17	—	0.003	0.009	0.010
		1988	137	—	0.008	0.004	0.160
		1990	2	—	0.000	0.000	0.003
		1996	2	—	0.000	0.000	0.003
<i>Mangifera parvifolia</i> Anacardiaceae	Recruits:	1986	232	—	0.091	0.093	0.106
		1988	15	—	0.009	0.003	0.008
		1990	37	—	0.006	0.035	0.005
		1996	44	—	0.009	0.039	0.005
<i>Mangifera pentandra</i> Anacardiaceae	Recruits:	1986	78	0.098	—	—	—
		1988	2	0.003	—	—	—
		1990	0	0.000	—	—	—
		1996	0	0.000	—	—	—
<i>Shorea beccariana</i> Dipterocarpaceae	Recruits:	1986	536	—	0.170	0.205	0.295
		1988	89	—	0.035	0.024	0.006
		1990	18	—	0.000	0.008	0.015
		1996	290	—	0.193	0.024	0.015
<i>Whiteodendron moultonianum</i> Myrtaceae	Recruits:	1986	359	—	0.329	0.018	0.103
		1988	72	—	0.048	0.000	0.043
		1990	15	—	0.008	0.001	0.010
		1996	33	—	0.025	0.000	0.010
Bako							
<i>Artocarpus odoratissimus</i> Moraceae	Recruits:	1986	21	0.005	0.003	0.007	0.014
		1988	13	0.005	0.001	0.004	0.007
		1990	4	0.000	0.004	0.001	0.000
		1996	1	0.000	0.000	0.001	0.000
<i>Cotylelobium melanoxylon</i> Dipterocarpaceae	Recruits:	1986	186	0.064	0.024	0.030	0.132
		1988	46	0.036	0.000	0.008	0.017
		1990	33	0.010	0.003	0.004	0.028
		1996	8	0.005	0.000	0.001	0.004
<i>Dryobalanops beccarii</i> Dipterocarpaceae	Recruits:	1986	458	0.051	0.284	—	0.289
		1988	41	0.009	0.007	—	0.040
		1990	17	0.001	0.012	—	0.010
		1996	1162	1.034	0.444	—	3.975
<i>Gonocaryum minus</i> Icacinaceae	Recruits:	1986	148	0.123	0.061	—	0.010
		1988	37	0.036	0.012	—	0.000
		1990	18	0.015	0.004	—	0.004
		1996	8	0.003	0.007	—	0.001
<i>Pimelodendron griffithianum</i> Euphorbiaceae	Recruits:	1986	20	0.001	0.003	0.012	0.011
		1988	8	0.001	0.005	0.000	0.004
		1990	3	0.000	0.000	0.001	0.003
		1996	0	0.000	0.000	0.000	0.000
<i>Swintonia schwenkii</i> Anacardiaceae	Recruits:	1986	480	—	—	0.395	0.249
		1988	88	—	—	0.070	0.043
		1990	18	—	—	0.014	0.010
		1996	201	—	—	0.158	0.113

At Lambir, 800 m² was sampled in each plot in every census. Transect areas sampled at Bako were as follows: Plot 1 = 780 m²; Plot 2 = 740 m²; Plot 3 = 760 m²; and Plot 4 = 720 m², with the exception of *D. beccarii* recruits in 1996 which were sampled in Plot 3 as above and subsampled in Plots 1 (580 m²), 2 (550 m²) and 4 (80 m²).

combined, the probability of survival by 10 y was 52% (Figure 1, Table 2). An additional 2962 recruits were tagged in 1988, 1990 and 1996. Fifty seedlings progressed to the sapling size-class (> 1 cm dbh) in the intervening 10 y, representing a promotion rate of around 1%. The proportion of seedlings present in 1986 that survived to 1996 was significantly greater in Lambir (57%) than at Bako (43%) ($\chi^2 = 64.7$, $P < 0.0001$) (Table 2). The proportions of seedlings surviving in each plot also differed significantly within each site (Lambir: $\chi^2 = 75.4$, $P < 0.001$; Bako: $\chi^2 = 36.3$, $P < 0.0001$). Likewise the proportions of seedlings of each species surviving differed significantly within each site (Lambir: $\chi^2 = 208$, $P < 0.0001$; Bako: $\chi^2 = 94.0$, $P < 0.0001$).

By 1996, large numbers of seedlings were at least a decade old, and probably many were much older. For seven of the 13 species (three at Bako and four at Lambir), 65–96% of seedlings present in 1996 were at least 10 y old (Figure 2).

Across sites, five of the 13 species had large seedling establishment events during the course of the 10 y study that were evident from the census work (Table 1). *Elateriospermum tapos* had a large number of new seedlings at the 1988 census. *Allantospermum borneensis*, *D. beccarii*, *S. beccariana* and *S. schwenkii* had large numbers of new seedlings at the 1996 census.

To determine if fast-growing seedlings continue to grow rapidly in successive time periods, and whether seedlings with poor growth performance can recover to grow rapidly again, we used census records of the five study species that had the largest population sizes at the initiation of the study: *Dipterocarpus globosus*, *D. beccarii*, *S. beccariana*, *S. schwenkii* and *W. moultonianum*. For each species, seedlings were ranked by their growth performance at each census. Six categories of seedling performance were described: (1) advanced to sapling: the seedling achieved 1 cm dbh; (2) rapid growth: annual growth rate (the 90th percentile of growth rates for seedlings of the same species in the census interval of interest); (3) moderate growth: annual height growth rate > 2 cm y^{-1} . (The measurement error for height was approximately ± 2 cm. The 2 cm cut-off was at the 50th percentile of positive growth for all species combined in all census intervals combined.); (4) suppressed: $|\text{change in height}| \leq 2$ cm y^{-1} ; (5) height loss: annual height loss > 2 cm y^{-1} ; (6) dead: seedling died during the census interval (Figure 3).

The fastest-growing seedlings from the 1986–1988 census interval were selected for an examination of their performance in 1988–1990. Seedlings were capable of surviving through periods of suppression or height loss after a period of fast growth (Figure 3). Of the fast-growing seedlings from the initial census interval, 25–60% showed poor performance in 1988–1990, but these seedlings survived to the end of the second census interval. Of this set of seedlings, a subset was selected that was composed of those initially fast-growing seedlings that were then suppressed or that lost height in 1988–1990. These seedlings were assessed for their performance in the 1990–1996 census interval. Of these fast-growing seedlings that had become suppressed or lost height, 21–63%




Figure 1. Survivorship of 13 shade-tolerant species in two mixed dipterocarp forests. Survival is shown as the percentage of seedlings surviving. Initial sample sizes are the same as those given in Table 1. Diamonds represent seedlings present in 1986. Squares represent seedlings that were new in 1988. Triangles represent seedlings that were new in 1990.

Table 2. Seedling performance over 10 y as shown by proportion surviving (%), and median and maximum annualized growth rates (cm y^{-1}) for seedlings present in 1986. N for growth rates reflects only surviving seedlings with positive height change.

		Survival		Growth rate		
		N	%	N	Median	Max
Lambir						
	Total	3308	56	1680	1.8	22.5
	Plot 2	78	33	24	0.9	7.5
	3	797	48	326	1.5	22.5
	4	1261	55	632	1.7	11.0
	5	1172	65	698	2.1	12.0
	<i>Allantospermum borneensis</i>	205	67	113	1.4	14.2
	<i>Dipterocarpsu globosus</i>	1881	61	1059	2.0	22.5
	<i>Elateriospermum tapos</i>	17	29	5	1.0	3.2
	<i>Mangifera parvifolia</i>	232	81	176	1.4	6.8
	<i>Mangifera pentandra</i>	78	33	24	0.9	7.5
	<i>Shorea beccariana</i>	536	43	184	2.0	12.0
	<i>Whiteodendron moultonianum</i>	359	37	119	1.0	15.8
Bako						
	Total	1314	43	433	2.0	20.0
	Plot 1	191	42	70	1.8	20.0
	2	277	65	59	1.6	7.5
	3	338	64	104	1.5	13.9
	4	507	53	200	2.5	15.9
	<i>Artocarpus odoratissimus</i>	21	48	8	2.9	9.4
	<i>Cotylelobium melanoxyton</i>	186	55	73	2.4	20.0
	<i>Dryobalanops beccarii</i>	458	38	129	3.0	15.3
	<i>Gonocaryum minus</i>	148	73	66	1.8	6.9
	<i>Pimelodendron griffithianum</i>	20	65	6	4.7	6.5
	<i>Swintonia schwenkii</i>	480	33	151	1.7	15.9

began to grow again in the third census interval, some moving into the sapling size-class, and as much as 15% rejoined the rapid-growth category.

DISCUSSION

Many tropical trees, particularly in South-East Asia, have supra-annual fruiting events (Ashton 1988, Corlett 1990, Yap 1987, Yasuda *et al.* 1999), and the newly germinated seedlings have a high mortality rate (Itoh 1995, Itoh *et al.* 1995, Turner 1990). However, this high mortality rate obscures the fact that those seedlings that do become established after the initial die-off of most of their cohort are resilient and long-lived. The long-term survival of the seedlings of shade-tolerant canopy tree species in tropical forests has also been documented in Australia (Connell & Green 2000, Connell *et al.* 1984), Panama (DeSteven 1994), French Guiana (Forget 1992, 1997) and Zaire (Hart 1995). For many tree species at Bako and Lambir, the majority of seedlings are a decade or more in age, and some seedlings may even be ≥ 50 y old.

In these forests, the seedling bank is a multi-aged stand composed of individuals from multiple fruiting events, in which a seedling's height is a poor




Figure 2. Per cent of all seedlings from each species that were at least 10 y old in 1996. The total number of seedlings of each species present in 1996 is given in parentheses.

predictor of its age. A small seedling that has never experienced a light gap might be from the same cohort of several decades ago as a canopy tree that germinated in a large gap and grew more quickly. The processes that take place in the understory community determine the set of seedlings present at the time and location of gap formation (Brokaw & Scheiner 1989), and thus are critical for the regeneration of mixed dipterocarp forests.

The tree seedlings in these populations that have the fastest initial growth rates can become suppressed and survive for months or years with slow growth. The ability of seedlings to persist and grow slowly in the shaded understory may confer a great advantage as height is a strong determinant of seedling survival and competitive ability at the time of gap formation (Brown & Whitmore 1992, Connell 1989). Furthermore, old, previously suppressed seedlings in these forests can recover from suppression to achieve short-term growth




Figure 3. Alternating performance across three consecutive census intervals of seedlings tagged in the initial 1986 census. Each bar represents the relative proportions of seedlings in each of six performance rankings in census intervals ending in 1988, 1990 and 1996. The dotted lines indicate the subset of seedlings in each census interval that were examined for performance rank in the subsequent census interval. Seedlings with the highest performance rank (Rapid growth) in the first census interval were followed into the second census interval. Of these, seedlings with the poor performance ranks (Height loss or Suppressed) were followed into the third census interval.

rates equivalent to those of the fastest-growing seedlings in the population. A similar plasticity of growth rates has been noted for dipterocarp seedlings and saplings in Sabah in northern Borneo, and in Sri Lanka (Ashton & de Zoysa 1990, Moad 1992). This capacity to survive through alternating periods of fast and slow growth facilitates the persistence of seedlings over time, allowing access to a greater number of small, temporary gaps, and so should be selectively advantageous if these seedlings have a chance of reaching maturity.

These results have implications for forestry methods that require an assessment of the vigour of pre-existing seedlings before harvest and that are based on the untested assumption that older individuals in low-light environments lose their ability to respond to improved growing conditions (Ashton 1990, Poore 1968, Tang 1987). Our study provides evidence that 10-y-old seedlings that have persisted through periods of poor performance can return to rapid growth, and supports the idea that nearly all seedlings can respond to an opening in the canopy regardless of age (Liew & Wong 1973).

Space for establishment, in this case canopy gaps, allocated at random and paired with environmental variability may facilitate species coexistence (Chesson & Warner 1981, Hubbell & Foster 1986, Sale 1977). The rarity and small size of canopy gaps in these forests may enhance the importance of recruitment limitation, in this case failure to have seedlings at the site of a gap, as a controlling factor in community diversity and composition (Brokaw & Busing 2000, Hubbell *et al.* 1999, Hurtt & Pacala 1995). The ability of seedlings to persist in a suppressed state under deep shade has been cited as a possible mechanism for the maintenance of high levels of diversity in the tropical rain forest (Uhl *et al.* 1988). This ability would allow weak competitors access to gaps inaccessible to stronger competitors that happen to be recruitment limited (Brown *et al.* 1999, Wright 2002). An adaptation for the long-term persistence of seedlings could help to explain the unusually high levels of species diversity in the relatively stable forests of north-west Borneo.

ACKNOWLEDGEMENTS

We gratefully acknowledge the Forest Department of Sarawak for their kind assistance with this project. In particular, we would like to acknowledge the great contribution of Sylvester Tan in the management of census work and data. We also thank Peter S. Ashton, Adrien Finzi, Thomas Kunz and Gillian Cooper-Driver, and three anonymous reviewers for their invaluable comments on various versions of this manuscript. This work was supported by NSF Grant DEB - 9974133, in-kind support from the Center for Tropical Forest Science of the Smithsonian Tropical Research Institute, and a grant from the Sarawak Timber Association.

LITERATURE CITED

- ADAMS, P. 1992. *Australian rainforests*. Clarendon Press, Oxford. 308 pp.
- APPANAH, S. 1985. General flowering in the climax rainforests of Southeast Asia. *Journal of Tropical Ecology* 1:225–240.
- ARRIAGA, L. 2000. Types and causes of tree mortality in a tropical montane cloud forest of Tamaulipas, Mexico. *Journal of Tropical Ecology* 16:623–636.
- ASHTON, P. M. S. 1990. Method for the evaluation of advanced regeneration. *Forest Ecology and Management* 36:163–175.
- ASHTON, P. M. S. & DE ZOYSA, N. D. 1990. Performance of *Shorea trapezifolia* (Thwaites) Ashton seedlings growing in different light regimes. *Journal of Tropical Forest Science* 1:358–364.
- ASHTON, P. S. 1969. Speciation among tropical forest trees: some deductions in the light of recent evidence. *Biological Journal of the Linnean Society* 1:155–196.
- ASHTON, P. S. 1988. Dipterocarp biology as a window to the understanding of tropical forest structure. *Annual Review of Ecology and Systematics* 19:347–370.
- ASHTON, P. S. & HALL, P. 1992. Comparisons of structure among mixed dipterocarp forests of north-western Borneo. *Journal of Ecology* 80:459–481.
- BAILLIE, I. C. 1976. Further studies on drought in Sarawak, East Malaysia. *Journal of Tropical Geography* 43:20–29.
- BARNARD, R. C. 1956. Recruitment, survival, and growth of timber-tree seedlings in natural rain forest. *The Malayan Forester* XIX:156–161.
- BEAMAN, R. S., BEAMAN, J. H., MARSH, C. W. & WOODS, P. V. 1985. Drought and forest fires in Sabah in 1983. *Sabah Society Journal* 8:10–30.
- BROKAW, N. & BUSING, R. T. 2000. Niche versus chance and tree diversity in gaps. *Trends in Ecology and Evolution* 15:183–188.
- BROKAW, N. V. L. & SCHEINER, S. M. 1989. Species composition in gaps and structure of a tropical rain forest. *Ecology* 70:538–541.
- BROKAW, N. V. L. & WALKER, L. R. 1991. Summary of the effects of Caribbean hurricanes on vegetation. *Biotropica* 23:442–447.
- BROWN, N. D. & WHITMORE, T. C. 1992. Do dipterocarp seedlings really partition tropical rain forest gaps? *Philosophical Transactions of the Royal Society of London, Series B* 335:369–378.
- BROWN, N., PRESS, M. & BEBBER, D. 1999. Growth and survivorship of dipterocarp seedlings: differences in shade persistence create a special case of dispersal limitation. *Philosophical Transactions: Biological Sciences* 354:1847–1855.
- BRÜNIC, E. F. 1969. On the seasonality of droughts in the lowlands of Sarawak (Borneo). *Erdkunde* 23:127–133.
- CHESSON, P. & WARNER, R. 1981. Environmental variability promotes coexistence in lottery competitive systems. *American Naturalist* 117:923–943.
- CLARK, D. B. 1990. The role of disturbance in the regeneration of neotropical moist forests. Pp. 291–315 in Bawa, K. & Hadley, M. (eds). *Reproductive ecology of tropical forest plants*. UNESCO, Paris and Parthenon Publishing, Carnforth. 421 pp.
- CONDIT, R. 1995. Research in large, long-term tropical forest plots. *Trends in Ecology and Evolution* 10:18–22.
- CONNELL, J. H. 1989. Some processes affecting the species composition in forest gaps. *Ecology* 70:560–562.
- CONNELL, J. H. & GREEN, P. T. 2000. Seedling dynamics over thirty-two years in a tropical rain forest tree. *Ecology* 81:568–584.
- CONNELL, J. H., TRACEY, J. G. & WEBB, L. J. 1984. Compensatory recruitment, growth, and mortality as factors maintaining rain forest tree diversity. *Ecological Monographs* 54:141–164.
- CORLETT, R. T. 1990. Flora and reproductive phenology of the rain forest at Bukit Timah, Singapore. *Journal of Tropical Ecology* 6:55–63.
- CROW, T. R. 1980. A rainforest chronicle: a 30-year record of change in structure and composition at El Verde, Puerto Rico. *Biotropica* 12:42–55.
- CURRAN, L. M., CANIAGO, I., PAOLI, G. D., ASTIANI, D., KUSNETI, M., LEIGHTON, M., NIRARITA, C. E. & HAERUMAN, H. 1999. Impact of El Niño and logging on canopy tree recruitment in Borneo. *Science* 286:2184–2188.
- CURRAN, L. M. & WEBB, C. O. 2000. Experimental tests of the spatiotemporal scale of seed predation in mast-fruited Dipterocarpaceae. *Ecological Monographs* 70:129–148.
- DAVIES, S. J., PALMIOTTO, P., ASHTON, P. S., LEE, H. S. & LAFRANKIE, J. V. 1998. Comparative ecology of 11 sympatric species of *Macaranga* in Borneo: tree distribution in relation to horizontal and vertical resource heterogeneity. *Journal of Ecology* 86:662–673.
- DELISSIO, L. J. 2000. *Tree seedling dynamics and drought in Bornean rain forests*. Ph.D. dissertation. Department of Biology, Boston University, Boston. 239 pp.

- DESTEVEN, D. 1994. Tropical tree seedling dynamics: recruitment patterns and their population consequences for three canopy species in Panama. *Journal of Tropical Ecology* 10:369–383.
- FORGET, P.-M. 1992. Regeneration ecology of *Eperua grandiflora* (Caesalpinaceae), a large seeded tree in French Guiana. *Biotropica* 24:146–156.
- FORGET, P.-M. 1997. Ten-year seedling dynamics in *Vouacoua americana* in French Guiana: a hypothesis. *Biotropica* 29:124–126.
- FOX, J. E. D. 1967. An enumeration of lowland dipterocarp forest in Sabah. *Malayan Forester* 30:263–279.
- GALE, N. 2000. The aftermath of tree death: coarse woody debris and the topography in four tropical rain forests. *Canadian Journal of Forest Research* 30:1489–1493.
- GARWOOD, N. C., JANOS, D. P. & BROKAW, N. 1979. Earthquake-caused landslides: a major disturbance to tropical forests. *Science* 205:997–999.
- HALL, P. 1991. *Structure, stand dynamics, and species compositional change in three mixed dipterocarp forests of northwest Borneo*. Ph.D. dissertation. Department of Biology, Boston University, Boston. 701 pp.
- HALL, P. & PRIMACK, R. B. 1987. *Long-term ecological studies in three contrasting mixed dipterocarp forests*. Boston University. Unpublished report. Boston University, Boston. 65 pp.
- HART, B. 1995. Seed, seedling and sub-canopy survival in monodominant and mixed forests of the Ituri Forest, Africa. *Journal of Tropical Ecology* 11:443–459.
- HARTSHORN, G. S. 1978. Tree falls and tropical forest dynamics. Pp. 617–630 in Tomlinson, P. B. & Zimmermann, M. H. (eds). *Tropical trees as living systems: the proceedings of the Fourth Cabot Symposium held at Harvard Forest, Petersham, Massachusetts, on April 26–30, 1978*. Cambridge University Press, Cambridge. 675 pp.
- HUBBELL, S. P. & FOSTER, R. B. 1986. Biology, chance and history and the structure of tropical rain forest tree communities. Pp. 314–329 in Diamond, J. M. & Case, T. J. (eds). *Community ecology*. Harper and Row, New York. 665 pp.
- HUBBELL, S. P., FOSTER, R. B., O'BRIEN, S. T., HARMS, K. E., CONDIT, R., WECHSLER, B., WRIGHT, S. J. & LAO, S. L. D. 1999. Light gap disturbances, recruitment limitation, and tree diversity in a neotropical forest. *Science* 283:554–557.
- HURTT, G. C. & PACALA, S. W. 1995. The consequences of recruitment limitation: reconciling chance, history and competitive differences between plants. *Journal of Theoretical Biology* 176:1–12.
- ITOH, A. 1995. Effects of forest floor environment on germination and seedling establishment of two Bornean rainforest emergent species. *Journal of Tropical Ecology* 11:517–527.
- ITOH, A., YAMAKURA, T., OGINO, K. & LEE, H. S. 1995. Survivorship and growth of seedlings of four dipterocarp species in a tropical rainforest of Sarawak, East Malaysia. *Ecological Research* 10:327–338.
- ITOH, A., YAMAKURA, T., OGINO, K., LEE, H. S. & ASHTON, P. 1997. Spatial distribution of two predominant emergent trees in a tropical rainforest in Sarawak, Malaysia. *Plant Ecology* 132:121–136.
- LEE, H. S., ASHTON, P. S. & OGINO, K. 1995. *Long term ecological research of tropical rain forest in Sarawak*. Ehime University, Matsuyama. 228 pp.
- LIEW, T. C. & WONG, F. O. 1973. Density, recruitment, mortality and growth of dipterocarp seedlings in virgin and logged-over forests in Sabah. *The Malaysian Forester* 36:3–15.
- MOAD, A. S. 1992. *Dipterocarp juvenile growth and understory light availability in Malaysian tropical forest*. Ph.D. dissertation. Harvard University, Cambridge, MA. 328 pp.
- NAKAGAWA, M., TANAKA, K., NAKASHIZUKA, T., OHKUBO, T., KATO, T., MAEDA, T., SATO, K., MIGUCHI, H., NAGAMASU, H., OGINO, K., TEO, S., HAMID, A. A. & LEE, H. S. 2000. Impact of severe drought associated with the 1997–1998 El Niño in a tropical forest in Sarawak. *Journal of Tropical Ecology* 16:355–367.
- OHKUBO, T., MAEDA, T., KATO, T., TANI, M., YAMAKURA, T., LEE, H. S., ASHTON, P. S. & OGINO, K. 1995. Landslide scars in canopy mosaic structure as a large scale disturbance to a mixed-dipterocarp forest at Lambir Hills National Park, Sarawak. Pp. 172–184 in Lee, H. S., Ashton, P. S. & Oginno, K. (eds). *Long term ecological research of tropical rain forest in Sarawak*. Ehime University, 228 pp.
- OHKUBO, T., TANI, M., NOGUCHI, H., YAMAKURA, T., ASHTON, P. S. & OGINO, K. Submitted. *Topographic pattern of canopy gap formation in a mixed dipterocarp forest in Northwestern Borneo*. International Meeting of the Center for Tropical Forest Science 1998, Washington, DC, Smithsonian Tropical Research Institute.
- PALMIOTTO, P. A. 1998. *The role of specialization in nutrient-use efficiency as a mechanism driving species diversity in a tropical rain forest*. Doctor of Forestry Dissertation. School of Forestry and Environmental Studies, Yale University, New Haven, Connecticut. 282 pp.
- PLATT, W. J. & STRONG, D. R. 1989. Gaps in forest ecology. *Ecology* 70:436–476.
- POORE, M. E. D. 1968. Studies in Malaysian rain forest: the forest on Triassic sediments in Jengka Forest Reserve. *Journal of Ecology* 56:143–196.

- PRIMACK, R. B. & HALL, P. 1992. Biodiversity and forest change in Malaysian Borneo. *BioScience* 42:829–837.
- PUTZ, F. E. & MILTON, K. 1982. Tree mortality rates on Barro Colorado Island. Pp. 95–100 in Leigh, E. G., Rand, A. S. & Windsor, D. M. (eds). *The ecology of a tropical forest: seasonal rhythms and long-term changes*. Smithsonian Institution Press, Washington, DC. 503 pp.
- SALE, P. F. 1977. Maintenance of high diversity in coral reef communities. *American Naturalist* 111:337–359.
- SWAINE, M. D. & WHITMORE, T. C. 1988. On the definition of ecological species groups in tropical rain forests. *Vegetatio* 75:81–86.
- TANG, H. T. 1987. Problems and strategies for regenerating dipterocarp forests in Malaysia. Pp. 24–41 in Merger, F. & Vincent, J. R. (eds). *Natural management of tropical moist forest*. Yale School of Forestry and Environmental Studies, New Haven.
- TURNER, I. M. 1990. Tree seedling growth and survival in a Malaysian rain forest. *Biotropica* 22:146–154.
- UHL, C., CLARK, K., DEZZEO, N. & MAQUIRINO, P. 1988. Vegetation dynamics in Amazon treefall gaps. *Ecology* 69:751–763.
- VAN DER MEER, P. J. 1995. *Canopy dynamics of a tropical rain forest in French Guiana*. Ph.D. dissertation. Wageningen Agricultural University, Wageningen, The Netherlands.
- VITOUSEK, P. M., WALKER, L. R., WHITEAKER, L. D. & MATSON, P. A. 1993. Nutrient limitation to plant growth during primary succession in Hawaii Volcanoes National Park. *Biogeochemistry* 23:197–215.
- WALKER, L. R. 1991. Tree damage and recovery from Hurricane Hugo in Luquillo Experimental Forest, Puerto Rico. *Biotropica* 23:379–385.
- WALSH, R. P. D. 1996. Drought frequency changes in Sabah and adjacent parts of northern Borneo since the late nineteenth century and possible implications for tropical rain forest dynamics. *Journal of Tropical Ecology* 12:385–407.
- WHITMORE, T. C. 1978. Gaps in the forest canopy. Pp. 639–655 in Tomlinson, P. B. & Zimmerman, M. H. (eds). *Tropical trees as living systems*. Cambridge University Press, Cambridge. 675 pp.
- WHITMORE, T. C. 1984. *Tropical rain forests of the Far East*. Clarendon Press, Oxford. 352 pp.
- WHITMORE, T. C. 1989. Changes over twenty-one years in the Kolombangara rain forests. *Journal of Ecology* 77:469–483.
- WHITMORE, T. C. 1995. Comparing Southeast Asian and other tropical rain forests. Pp. 5–15 in Primack, R. B. & Lovejoy, T. E. (eds). *Ecology, conservation, and management of Southeast Asian rainforests*. Yale University Press, New Haven. 304 pp.
- WHITMORE, T. C. 1998. Potential impact of climate change on tropical rain forest seedlings and forest regeneration. Pp. 429–438 in Markham, A. (ed.). *Potential impacts of climate change on tropical forest ecosystems*. Kluwer Academic Publishers, Dordrecht. 463 pp.
- WICH, S. A. & VAN SCHAİK, C. P. 2000. The impact of El Niño on mast fruiting in Sumatra and elsewhere in Malesia. *Journal of Tropical Ecology* 16:563–577.
- WOOD, G. H. S. 1956. The dipterocarp flowering season in north Borneo, 1955. *Malayan Forester* 19:193–201.
- WRIGHT, S. J. 2002. Plant diversity in tropical forests: a review of species coexistence. *Oecologia* 130:1–14.
- WYATT-SMITH, J. 1958. Seedling/sapling survival of *Shorea leprosula*, *Shorea parvifolia*, and *Koompassia malaccensis*. *Malayan Forester* 21:185–193.
- YAMAKURA, T., KANZAKI, M., ITOH, A., OHKUBO, T., OGINO, K., CHAI, E. O. K., LEE, H. S. & ASHTON, P. S. 1995. Forest architecture of Lambir rain forest revealed by a large-scale research plot I. Topography of the plot. Pp. 2–20 in Lee, H. S., Ashton, P. S. & Oginno, K. (eds). *Long term ecological research of tropical rain forest in Sarawak*. Ehime University, Matsuyama. 228 pp.
- YAP, S. K. 1987. Gregarious flowering of dipterocarps: observations based on fixed tree populations in Selangor and Negri Sembilan, Malay Peninsula. Pp. 305–317 in Kostermans, A. J. G. H. (ed.). *Proceedings of the Third Round Table Conference on Dipterocarps*. UNESCO, Bogor. 657 pp.
- YASUDA, M., MATSUMOTO, J., OSADA, N., ICHIKAWA, S., KACHI, N., TANI, M., OKUDA, T., FURUKAWA, A., NIK, A. R. & MANOKARAN, N. 1999. The mechanism of general flowering in Dipterocarpaceae in the Malay Peninsula. *Journal of Tropical Ecology* 15:437–449.